









# sigvaris

## MEDICAL COMPRESSION PRESCRIPTION PAD

Patient	Date
Diagnosis	Number of refills
	Number of pairs
	No substitution

Physician Name	
Physician Signature 	
License Number	Phone

Compression (mmHg)				
<input type="checkbox"/> 15-20	<input type="checkbox"/> 20-30	<input type="checkbox"/> 30-40	<input type="checkbox"/> 40-50	<input type="checkbox"/> 50-60

<input type="checkbox"/> Closed Toe <input type="checkbox"/> Open Toe <input type="checkbox"/> Patient Choice						
						
<input type="checkbox"/> Calf <input type="checkbox"/> w/Grip-top	<input type="checkbox"/> Thigh	<input type="checkbox"/> Pantyhose	<input type="checkbox"/> L <input type="checkbox"/> R Thigh w/ waist	<input type="checkbox"/> Maternity/ Plus Sizes	<input type="checkbox"/> Wraps <input type="checkbox"/> w/Foot piece	<input type="checkbox"/> Armsleeve <input type="checkbox"/> w/Glove <input type="checkbox"/> w/Gauntlet

### Donning Devices:

Please see reverse side for indications

<input type="radio"/> Rubber Gloves	<input type="radio"/> SIMON	<input type="radio"/> Magnide®
<input type="radio"/> Latex-free Gloves	<input type="radio"/> Sim-Slide®	<input type="radio"/> SIGVARIS Doff'n Donner
<input type="radio"/> Textile Gloves	<input type="radio"/> Cotton Underliner	<input type="radio"/> SIGVARIS Doff'n Donner Cone



WEST END MAMAS  
*total wellness for motherhood*

# GRADUATED COMPRESSION CHART

	Symptoms/Indications	Compression strength at the ankle (mmHg)
<b>CEAP 0/A:</b>	<ul style="list-style-type: none"> <li>• Extended standing or sitting (travel)</li> <li>• Prophylaxis (pregnancy, risk factors)</li> </ul>	<b>15-20 mmHg</b>
<b>CEAP 1/S:</b>	<ul style="list-style-type: none"> <li>• Heavy, fatigued, tired legs</li> </ul>	
<b>CEAP 1/S:</b>	<ul style="list-style-type: none"> <li>• Painful, fatigued and aching legs</li> <li>• Spider veins/mild edema (due to pregnancy, age, travelling, etc.)</li> <li>• Post sclerotherapy of small veins</li> </ul>	<b>20-30 mmHg</b>
<b>CEAP 2/S:</b>	<ul style="list-style-type: none"> <li>• Mild to moderate varicose veins</li> <li>• Elective surgery (sclerotherapy, phlebectomy, vein stripping, orthopaedics)</li> <li>• Orthostatis/Postural hypotension</li> </ul>	
<b>CEAP 3/S:</b>	<ul style="list-style-type: none"> <li>• Moderate/severe edema (pregnancy, risk factors)</li> <li>• Surgery (orthopaedics, post fracture, traumatic edema, sclerotherapy, phlebectomy, vein stripping)</li> </ul>	<b>30-40 mmHg</b>
<b>CEAP 4/S:</b>	<ul style="list-style-type: none"> <li>• Skin changes without ulceration</li> <li>• Superficial Phlebitis (red, painful)</li> <li>• Deep Vein Thrombosis/ Post Thrombotic Syndrome</li> </ul>	
<b>CEAP 5/S:</b>	<ul style="list-style-type: none"> <li>• Skin changes with healed ulceration</li> </ul>	
<b>CEAP 6/S:</b>	<ul style="list-style-type: none"> <li>• Skin changes with active ulceration</li> </ul>	
<b>CEAP 4/S:</b>	<ul style="list-style-type: none"> <li>• Severe Deep Vein Thrombosis/Post Thrombotic Syndrome</li> </ul>	<b>40-50 or 50-60 mmHg</b>
<b>CEAP 5/S:</b>	<ul style="list-style-type: none"> <li>• Severe skin changes with healed ulceration (recurrent)</li> </ul>	
<b>CEAP 6/S:</b>	<ul style="list-style-type: none"> <li>• Severe skin changes with active ulceration</li> </ul>	

CONTRAINDICATIONS	CAUTION
<ul style="list-style-type: none"> <li>• Arterial insufficiency, intermittent claudication, ischemia</li> <li>• Uncontrolled congestive heart failure</li> <li>• Acute dermatitis, weeping dermatosis, cutaneous sepsis</li> </ul>	<ul style="list-style-type: none"> <li>• Signs of infection</li> <li>• Extensive venous ulceration</li> <li>• Skin sensitivities or allergies</li> <li>• Neuropathy</li> <li>• History of diabetes</li> <li>• Confinement to bed or non-ambulatory use unless otherwise prescribed by the physician</li> </ul>

No liability accepted for non-observance of contraindications and cautions.